

# Exam on Quantum Field Theory I

Devised by Timo Weigand, written on 03.04.2014

Solutions by Janosh Riebesell, 02.04.2015

## Problem 1 (11 points)

The action for a complex scalar field  $\varphi(x)$  is given by

$$S = \int d^4x (\partial_\mu \varphi^\dagger(x) \partial^\mu \varphi(x) - M^2 \varphi^\dagger(x) \varphi(x)). \quad (1)$$

The mode expansion of  $\varphi(x)$  is given by

$$\varphi(x) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} (a(p)e^{-ipx} + b^\dagger(p)e^{ipx}) \Big|_{p^0=E_p},$$

where the creation and annihilation operators satisfy the commutation relations

$$[a(p), a^\dagger(q)] = [b(p), b^\dagger(q)] = (2\pi)^3 \delta^{(3)}(\vec{p} - \vec{q}).$$

a) For a theory of several fields  $\chi_k$  the canonical stress-energy tensor is

$$T^{\mu\nu} = \sum_k \frac{\partial \mathcal{L}}{\partial(\partial_\mu \chi_k)} \partial^\nu \chi_k - \eta^{\mu\nu} \mathcal{L}.$$

Show that  $\partial_\mu T^{\mu\nu} = 0$  on shell for the theory of a complex scalar field given by (1).

$$\begin{aligned} T^{\mu\nu} &= \partial^\mu \varphi^\dagger \partial^\nu \varphi + \partial^\mu \varphi \partial^\nu \varphi^\dagger - \eta^{\mu\nu} \partial_\alpha \varphi^\dagger \partial^\alpha \varphi + \eta^{\mu\nu} M^2 \varphi^\dagger \varphi, \\ \partial_\mu T^{\mu\nu} &= \partial_\mu (\partial^\mu \varphi^\dagger \partial^\nu \varphi) + \partial_\mu (\partial^\mu \varphi \partial^\nu \varphi^\dagger) - \partial^\nu (\partial_\alpha \varphi^\dagger \partial^\alpha \varphi) + \partial^\nu (M^2 \varphi^\dagger \varphi) \\ &= \partial^2 \varphi^\dagger \partial^\nu \varphi + \underbrace{\partial_\mu \varphi^\dagger \partial_\mu \partial^\nu \varphi}_{=0 \text{ (e.o.m.)}} + \partial^2 \varphi \partial^\nu \varphi^\dagger + \underbrace{\partial_\mu \varphi \partial_\mu \partial^\nu \varphi^\dagger}_{=0 \text{ (e.o.m.)}} \\ &\quad - \underbrace{\partial_\nu \partial_\alpha \varphi^\dagger \partial^\alpha \varphi}_{=0 \text{ (e.o.m.)}} - \underbrace{\partial_\alpha \varphi^\dagger \partial^\nu \partial^\alpha \varphi}_{=0 \text{ (e.o.m.)}} + M^2 \varphi^\dagger \partial^\nu \varphi + M^2 \varphi \partial^\nu \varphi^\dagger \\ &= \underbrace{(\partial^2 \varphi^\dagger + M^2 \varphi^\dagger)}_{=0 \text{ (e.o.m.)}} \partial^\nu \varphi + \underbrace{(\partial^2 \varphi + M^2 \varphi)}_{=0 \text{ (e.o.m.)}} \partial^\nu \varphi^\dagger = 0. \end{aligned}$$

We see that on-shell, i.e. when imposing the Klein-Gordon eq. – the equation of motion for the free scalar field –  $(\partial^2 + M^2)\varphi = 0$  to be true, the stress-energy tensor's total derivative vanishes.

b) From the stress-tensor components we define conserved charges as  $Q^\nu = \int d^3x T^{0\nu}$ . Show that the charges  $Q^i$  for  $i \in \{1, 2, 3\}$  are given by

$$Q^i = \int \frac{d^3p}{(2\pi)^3} p^i (a^\dagger(p)a(p) + b^\dagger(p)b(p)).$$

*Hint:* Perform the integral  $\int d^3x$  before performing any momentum integral.

From part a) we can deduce  $T^{0i}$  with  $i \in \{1, 2, 3\}$  to be

$$T^{0i} = \partial^0 \varphi^\dagger \partial^i \varphi + \partial^0 \varphi \partial^i \varphi^\dagger = \partial_t \varphi^\dagger \frac{\partial}{\partial x_i} \varphi + \partial_t \varphi \frac{\partial}{\partial x_i} \varphi^\dagger,$$

where

$$\begin{aligned}\partial_t \varphi &= i \int \frac{d^3 p}{(2\pi)^3} \sqrt{\frac{E_p}{2}} (b^\dagger(p) e^{ipx} - a(p) e^{-ipx}), \\ \partial^i \varphi &= -i \int \frac{d^3 p}{(2\pi)^3} \frac{p^i}{\sqrt{2E_p}} (a(p) e^{-ipx} - b^\dagger(p) e^{ipx}).\end{aligned}$$

Inserting the above expressions into the charge, we get

$$\begin{aligned}Q^i &= \int d^3 x (\partial_t \varphi^\dagger \partial^i \varphi + \partial_t \varphi \partial^i \varphi^\dagger) = \frac{1}{2} \int \frac{d^3 p}{(2\pi)^3} p^i (a^\dagger(p) a(p) - a^\dagger(p) b^\dagger(-p) \\ &\quad - b(p) a(-p) + b(p) b^\dagger(p) + b^\dagger(p) b(p) - b^\dagger(p) a^\dagger(-p) - a(p) b(-p) + a(p) a^\dagger(p)) \\ &= \frac{1}{2} \int \frac{d^3 p}{(2\pi)^3} p^i (2a^\dagger(p) a(p) + 2b^\dagger(p) b(p) + 2 \cdot (2\pi)^3 \delta^{(3)}(0)) \\ &\quad - \frac{1}{2} \int \frac{d^3 p}{(2\pi)^3} \left( \underbrace{p^i (a^\dagger(p) b^\dagger(-p) + a^\dagger(-p) b^\dagger(p))}_{\text{antisymmetric under } p^i \rightarrow -p^i} + \underbrace{p^i (a(p) b(-p) + a(-p) b(p))}_{\text{antisymmetric under } p^i \rightarrow -p^i} \right) \\ &= \int \frac{d^3 p}{(2\pi)^3} p^i (a^\dagger(p) a(p) + b^\dagger(p) b(p)) + \int d^3 p p^i \delta^{(3)}(0).\end{aligned}$$

We could get an overall minus sign in front of the charge, which would be no issue since we allow charges to take positive and negative values and which we call which is up to us. The last term is formally divergent ( $\delta^{(3)}(0)$  is the volume of space) but its integrand is odd in  $p^i$ , so it vanishes under symmetric integration. (Unlike for the energy  $Q^0$ , the vacuum carries no momentum.)

**c)** Consider the action of  $Q^i$  on the states  $a^\dagger(k) |0\rangle$  and  $b^\dagger(q) |0\rangle$  and hence give a physical interpretation for the charges  $Q^i$ .

$$\begin{aligned}Q^i a^\dagger(k) |0\rangle &= \int \frac{d^3 p}{(2\pi)^3} p^i (a^\dagger(p) a(p) + b^\dagger(p) b(p)) a^\dagger(k) |0\rangle \\ &= \int \frac{d^3 p}{(2\pi)^3} p^i (a^\dagger(p) (a^\dagger(k) a(p) + (2\pi)^3 \delta^{(3)}(p - k)) + a^\dagger(k) b^\dagger(p) b(p)) |0\rangle \\ &= \int \frac{d^3 p}{(2\pi)^3} p^i a^\dagger(p) (2\pi)^3 \delta^{(3)}(p - k) |0\rangle = k^i a^\dagger(k) |0\rangle, \\ Q^i b^\dagger(q) |0\rangle &= \int \frac{d^3 p}{(2\pi)^3} p^i (a^\dagger(p) a(p) + b^\dagger(p) b(p)) b^\dagger(q) |0\rangle \\ &= \int \frac{d^3 p}{(2\pi)^3} p^i (b^\dagger(q) a^\dagger(p) a(p) + b^\dagger(p) (b^\dagger(q) b(p) + (2\pi)^3 \delta^{(3)}(p - q))) |0\rangle \\ &= \int \frac{d^3 p}{(2\pi)^3} p^i b^\dagger(p) (2\pi)^3 \delta^{(3)}(p - q) |0\rangle = q^i b^\dagger(q) |0\rangle,\end{aligned}$$

where we used that the annihilation operators  $a(p)$ ,  $b(p)$  destroy the vacuum

$$a(p) |0\rangle = b(p) |0\rangle = 0 \quad \forall p.$$

We find that  $Q^i$ 's eigenvalues corresponding to the 1-particle states  $a^\dagger(k) |0\rangle$  and  $b^\dagger(q) |0\rangle$  are the spatial components of the momenta  $k$  and  $q$ .

## Problem 2 (6 points)

The Lagrangian density for a massless vector field  $A^\mu$  is given by

$$\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu}, \quad \text{where } F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu. \quad (2)$$

a) What problem does arise if one tries to quantise the theory given by eq. (2) by enforcing canonical equal-time commutation relations

$$[A^\mu(t, \vec{x}), \pi_\nu(t, \vec{y})] = i\delta_\nu^\mu \delta^{(3)}(\vec{x} - \vec{y})?$$

The Lagrangian in eq. (2) is not suitable for quantisation because it produces an unphysical conjugate momentum density. Specifically,  $\pi_\mu$  turns out to be

$$\pi_\mu = \frac{\partial \mathcal{L}}{\partial \dot{A}^\mu} = F_{\mu 0},$$

which yields immediately that  $\pi_0 = F_{00} = 0$  since  $F_{\mu\nu}$  is antisymmetric. Hence we must concede that  $(A^\mu, \pi_\mu)$  are no good canonical variables.

b) Sketch briefly how the problem can be overcome.

The problem stems from the gauge redundancy of  $A^\mu$ : not all four components are physical degrees of freedom, and  $A^0$  has no conjugate momentum. It can be overcome by fixing the gauge, e.g. by adding the Lorenz gauge-fixing term  $-\frac{1}{2\xi} (\partial_\mu A^\mu)^2$  to the Lagrangian. This yields  $\pi_0 = -\frac{1}{\xi} \partial_\mu A^\mu \neq 0$ , so the canonical commutation relations can be imposed on all four components. The unphysical (timelike and longitudinal) photon states this introduces are removed from the physical Hilbert space by the Gupta-Bleuler condition  $\partial_\mu A^{\mu,+} |\text{phys}\rangle = 0$ , leaving all physical observables unchanged.

## Problem 3 (12 points)

The Lagrangian density for a free Dirac fermion is given by

$$\mathcal{L}(x) = \bar{\Psi}(x)(i\gamma^\mu \partial_\mu - m)\Psi(x). \quad (3)$$

Consider a Lorentz transformation where we keep the coordinate axes fixed and only transform the fields, i.e.

$$x^\mu \rightarrow x^\mu \text{ (unchanged)}, \quad \partial^\mu \rightarrow \partial^\mu \text{ (unchanged)}, \quad \Psi(x) \rightarrow \Lambda_{\frac{1}{2}} \Psi(\Lambda^{-1}x),$$

where  $\Lambda_{\frac{1}{2}} = \exp(\frac{i}{2}\omega_{\mu\nu}S^{\mu\nu})$  with  $S^{\mu\nu} = \frac{i}{4}[\gamma^\mu, \gamma^\nu]$ . We also work in a basis where

$$\gamma^0 = \begin{pmatrix} 0 & \mathbb{1}_2 \\ \mathbb{1}_2 & 0 \end{pmatrix}, \quad \gamma^i = \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}$$

for  $i \in \{1, 2, 3\}$ , where  $\mathbb{1}_2$  is the  $2 \times 2$  unit matrix and  $\sigma^i$  are the Pauli matrices.

a) Show that  $(S^{\mu\nu})^\dagger \gamma^0 = \gamma^0 S^{\mu\nu}$ .

$$(S^{\mu\nu})^\dagger \gamma^0 = -\frac{i}{4} [(\gamma^\nu)^\dagger, (\gamma^\mu)^\dagger] \gamma^0 = \frac{i}{4} ((\gamma^\mu)^\dagger (\gamma^\nu)^\dagger \gamma^0 - (\gamma^\nu)^\dagger (\gamma^\mu)^\dagger \gamma^0)$$

$$= \frac{i}{4} \left( \underbrace{\gamma^0 \gamma^0}_{\mathbb{1}_4} (\gamma^\mu)^\dagger \gamma^0 \gamma^0 (\gamma^\nu)^\dagger \gamma^0 - \gamma^0 \gamma^0 (\gamma^\nu)^\dagger \gamma^0 \underbrace{\gamma^0 (\gamma^\mu)^\dagger \gamma^0}_{\gamma^\mu} \right) = \gamma^0 \frac{i}{4} [\gamma^\mu, \gamma^\nu] = \gamma^0 S^{\mu\nu},$$

where we used that  $(\gamma^0)^2 = \mathbb{1}_4$ , which follows directly from the defining relation of the Clifford algebra  $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu} \mathbb{1}_4$  for  $\mu = \nu = 0$ , and that  $(\gamma^\mu)^\dagger = \gamma^0 \gamma^\mu \gamma^0$  which holds specifically for our choice of gamma matrices, where  $(\gamma^0)^\dagger = \gamma^0$  and  $(\gamma^i)^\dagger = -\gamma^i$ . Together with the Clifford algebra's anticommutator, this yields

$$(\gamma^\mu)^\dagger = \begin{cases} \gamma^\mu = \gamma^0 \gamma^0 \gamma^\mu = \gamma^0 \gamma^\mu \gamma^0 & \text{for } \mu = 0 \\ -\gamma^\mu = -\gamma^\mu \gamma^0 \gamma^0 = \gamma^0 \gamma^\mu \gamma^0 & \text{for } \mu \in \{1, 2, 3\}. \end{cases}$$

**b)** Using the result from part a) and the fact that  $(\Lambda_{\frac{1}{2}})^{-1} \gamma^\mu \Lambda_{\frac{1}{2}} = \Lambda_\nu^\mu \gamma^\nu$  show that under a Lorentz transformation the Lagrangian density transforms as  $\mathcal{L}(x) \rightarrow \mathcal{L}(\Lambda^{-1}x)$ .

$$\begin{aligned} \mathcal{L}(x) &\xrightarrow{\Lambda_{\frac{1}{2}}} (\Lambda_{\frac{1}{2}} \Psi(\Lambda^{-1}x))^\dagger \gamma^0 (i\gamma^\mu \partial_\mu - m) \Lambda_{\frac{1}{2}} \Psi(\Lambda^{-1}x) \\ &= \Psi^\dagger(\Lambda^{-1}x) e^{-\frac{i}{2} \omega_{\mu\nu} (S^{\mu\nu})^\dagger} \gamma^0 (i\gamma^\mu \partial_\mu - m) \Lambda_{\frac{1}{2}} \Psi(\Lambda^{-1}x) \\ &= \bar{\Psi}(\Lambda^{-1}x) \underbrace{e^{-\frac{i}{2} \omega_{\mu\nu} S^{\mu\nu}}}_{(\Lambda_{\frac{1}{2}})^{-1}} (i\gamma^\mu \partial_\mu - m) \Lambda_{\frac{1}{2}} \Psi(\Lambda^{-1}x) \\ &= \bar{\Psi}(\Lambda^{-1}x) (i\Lambda_\nu^\mu \gamma^\nu \partial_\mu - m) \Psi(\Lambda^{-1}x) = \mathcal{L}(\Lambda^{-1}x), \end{aligned}$$

where we used that

$$e^{(S^{\mu\nu})^\dagger} \gamma^0 = \sum_{n=0}^{\infty} \frac{(S^{\mu\nu})^{\dagger n}}{n!} \gamma^0 \stackrel{\text{a)}}{=} \gamma^0 \sum_{n=0}^{\infty} \frac{(S^{\mu\nu})^n}{n!} = \gamma^0 e^{S^{\mu\nu}}$$

and in the last step the chain rule  $\partial_\mu(\Psi(\Lambda^{-1}x)) = [\Lambda^{-1}]_\mu^\rho (\partial_\rho \Psi)(\Lambda^{-1}x)$  together with  $\Lambda_\nu^\mu [\Lambda^{-1}]_\mu^\rho = \delta_\nu^\rho$ .

**c)** Define  $\Psi_\pm = P_\pm \Psi$  where  $P_\pm = \frac{1}{2}(\mathbb{1}_4 \pm \gamma^5)$  and  $\gamma^5 = i\gamma^0 \gamma^1 \gamma^2 \gamma^3$ . Show that  $\bar{\Psi}_\pm = \bar{\Psi} P_\mp$ . Rewrite the Lagrangian density (3) in terms of  $\Psi_+$  and  $\Psi_-$ . Derive the equation of motion for  $\Psi_+$  and  $\Psi_-$ . *Note:* You can use identities involving  $\gamma^5$  without proof.

$$\begin{aligned} (\gamma^5)^\dagger &= -i(\gamma^3)^\dagger (\gamma^2)^\dagger (\gamma^1)^\dagger (\gamma^0)^\dagger = -i(-\gamma^3)(-\gamma^2)(-\gamma^1)\gamma^0 = i\gamma^3 \gamma^2 \gamma^1 \gamma^0 = (-1)^3 i\gamma^0 \gamma^3 \gamma^2 \gamma^1 \\ &= (-1)^5 i\gamma^0 \gamma^1 \gamma^3 \gamma^2 = (-1)^6 i\gamma^0 \gamma^1 \gamma^2 \gamma^3 = \gamma^5, \\ \bar{\Psi}_\pm &= (\Psi_\pm)^\dagger \gamma^0 = (P_\pm \Psi)^\dagger \gamma^0 = \Psi^\dagger P_\pm^\dagger \gamma^0 = \Psi^\dagger \frac{1}{2} (\mathbb{1}_4 \pm (\gamma^5)^\dagger) \gamma^0 = \Psi^\dagger \gamma^0 \frac{1}{2} (\mathbb{1}_4 \pm (-1)^3 \gamma^5) \\ &= \bar{\Psi} \frac{1}{2} (\mathbb{1}_4 \mp \gamma^5) = \bar{\Psi} P_\mp. \end{aligned}$$

To rewrite the Lagrangian in eq. (3) in terms of  $\Psi_+$  and  $\Psi_-$ , we note that

$$\begin{aligned} P_+ + P_- &= \frac{1}{2}(\mathbb{1}_4 + \gamma^5) + \frac{1}{2}(\mathbb{1}_4 - \gamma^5) = \mathbb{1}_4 \implies \Psi_+ + \Psi_- = P_+ \Psi + P_- \Psi = (P_+ + P_-) \Psi = \Psi, \\ \bar{\Psi} &= \Psi^\dagger \gamma^0 = (\Psi_+ + \Psi_-)^\dagger \gamma^0 = \Psi_+^\dagger \gamma^0 + \Psi_-^\dagger \gamma^0 = \bar{\Psi}_+ + \bar{\Psi}_-, \\ \mathcal{L} &= \bar{\Psi}_+ (i\gamma^\mu \partial_\mu - m) \Psi_+ + \bar{\Psi}_+ (i\gamma^\mu \partial_\mu - m) \Psi_- + \bar{\Psi}_- (i\gamma^\mu \partial_\mu - m) \Psi_+ + \bar{\Psi}_- (i\gamma^\mu \partial_\mu - m) \Psi_- \\ &= \bar{\Psi}_+ i\gamma^\mu \partial_\mu \Psi_+ - \bar{\Psi}_+ m \Psi_+ + \bar{\Psi}_+ i\gamma^\mu \partial_\mu \Psi_- - \bar{\Psi}_+ m \Psi_- \\ &\quad + \bar{\Psi}_- i\gamma^\mu \partial_\mu \Psi_+ - \bar{\Psi}_- m \Psi_+ + \bar{\Psi}_- i\gamma^\mu \partial_\mu \Psi_- - \bar{\Psi}_- m \Psi_- \end{aligned}$$

$$\begin{aligned}
& +\bar{\Psi}_- i\gamma^\mu \partial_\mu \Psi_+ - \bar{\Psi}_- m \Psi_+ + \bar{\Psi}_- i\gamma^\mu \partial_\mu \Psi_- - \bar{\Psi}_- m \Psi_- \\
& = \bar{\Psi}_+ i\gamma^\mu \partial_\mu \Psi_+ + \bar{\Psi}_- i\gamma^\mu \partial_\mu \Psi_- - \bar{\Psi}_+ m \Psi_- - \bar{\Psi}_- m \Psi_+,
\end{aligned}$$

where we used in the last step that  $\bar{\Psi}_\pm \Psi_\pm = \bar{\Psi}_\mp P_\pm \Psi_\pm = 0$  and  $\bar{\Psi}_\pm \gamma^\mu \Psi_\mp = \bar{\Psi}_\mp P_\mp \gamma^\mu \Psi_\mp = \bar{\Psi}_\mp \gamma^\mu P_\pm \Psi_\mp = 0$  because  $P_\pm P_\mp = \frac{1}{4}(\mathbb{1}_4 - (\gamma^5)^2) = \frac{1}{4}(\mathbb{1}_4 - \mathbb{1}_4) = 0$ . Applying the Euler-Lagrange equation  $\partial_\mu \frac{\partial \mathcal{L}}{\partial(\partial_\mu \Psi)} = \frac{\partial \mathcal{L}}{\partial \Psi}$  to the new Lagrangian gives

$$\begin{aligned}
\Psi_+ : \quad \partial_\mu \bar{\Psi}_+ i\gamma^\mu + m \bar{\Psi}_- &= 0, \quad \bar{\Psi}_+ : \quad i\gamma^\mu \partial_\mu \Psi_+ - m \Psi_- = 0, \\
\Psi_- : \quad \partial_\mu \bar{\Psi}_- i\gamma^\mu + m \bar{\Psi}_+ &= 0, \quad \bar{\Psi}_- : \quad i\gamma^\mu \partial_\mu \Psi_- - m \Psi_+ = 0.
\end{aligned}$$

**d)** The spinors  $\Psi_+$  and  $\Psi_-$  correspond to two-component Weyl spinors. What are the properties of a particle which can be represented by a single Weyl spinor?

For a particle to be described by a single Weyl spinor, the above equations of motion would have to be decoupled. We can see that this is only the case for  $m = 0$ . Such a particle is massless and has definite chirality, which for a massless particle coincides with its helicity  $\pm \frac{1}{2}$ .

#### Problem 4 (11 points)

Consider the following theory for a complex scalar field  $\varphi$  and a real scalar field  $\sigma$ :

$$\mathcal{L} = \partial_\mu \varphi^\dagger \partial^\mu \varphi + \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma - M^2 \varphi^\dagger \varphi - \frac{m^2}{2} \sigma^2 - \lambda \sigma \varphi^\dagger \varphi - \lambda^2 \sigma^2 \varphi^\dagger \varphi.$$

**a)** State the momentum space Feynman rules for calculating scattering amplitudes for this theory. (No calculation required.)


To compute the amplitude for a scattering process with  $n$  ingoing momenta  $p_1, \dots, p_n$  and  $r$  outgoing momenta  $q_1, \dots, q_r$  in this theory, we perform the following:

- Draw all relevant fully connected amputated diagrams with  $(n + r)$  external points to given order in  $\lambda$ .
- Assign ingoing momenta labels  $p_k$ ,  $k \in \{1, \dots, n\}$  and outgoing momenta labels  $q_l$ ,  $l \in \{1, \dots, r\}$  and label momenta of internal lines with  $k_j$ .
- Each three-particle vertex gets a factor  $-i\lambda$ , each four-particle vertex  $-2i\lambda^2$  (the factor 2 from the two identical  $\sigma$  legs).
- To each internal line, we associate a propagator

$$\begin{aligned}
& \frac{i}{k_j^2 - M^2} \quad \text{if it is a } \varphi \text{ field,} \\
& \frac{i}{k_j^2 - m^2} \quad \text{if it is a } \sigma \text{ field.}
\end{aligned}$$

- We then integrate over all undetermined internal momenta  $\prod_j \int \frac{d^4 k_j}{(2\pi)^4}$ .
- Sum up all diagrams divided by their symmetry factors and multiply by  $\sqrt{Z}^{n+r}$  to given order in  $\lambda$ .

**b)** Draw all Feynman graphs to order  $\lambda^2$  for the non-trivial scattering of  $\varphi^\dagger \varphi \rightarrow \sigma \sigma$ .




c) Now write down the matrix element  $i\mathcal{M}$  for the process  $\varphi^\dagger\varphi \rightarrow \sigma\sigma$  to order  $\lambda^2$ . What is the matrix element  $i\mathcal{M}$  for the process  $\varphi\varphi \rightarrow \sigma\sigma$  to order  $\lambda^2$ ?

$$i\mathcal{M} = -2i\lambda^2 + (-i\lambda)^2 \left( \frac{i}{(p_1 - q_1)^2 - M^2} + \frac{i}{(p_1 - q_2)^2 - M^2} \right) = -i\lambda^2 \left( 2 + \frac{1}{(p_1 - q_1)^2 - M^2} + \frac{1}{(p_1 - q_2)^2 - M^2} \right).$$

The process  $\varphi\varphi \rightarrow \sigma\sigma$  violates the conservation of the  $U(1)$  charge (both interaction terms contain  $\varphi^\dagger\varphi$ ), so its amplitude vanishes,  $i\mathcal{M} = 0$ , at order  $\lambda^2$  and indeed to all orders.

d) Draw Feynman diagrams to and including order  $\lambda^4$  which contribute to the 1PI self energy (i.e. the 1PI contributions to the propagator) of the field  $\sigma$ .



The first two diagrams are the complete one-loop contributions (both of order  $\lambda^2$ ). At order  $\lambda^4$  there are a number of two-loop diagrams, the third being one example; the others are obtained by inserting a further cubic or quartic vertex into the  $\varphi$  loops of the first two diagrams.

## Problem 5 (9 points)

a) Discuss the difference between the bare mass  $m_0$  and the physical mass  $m$  of an electron in Quantum Electrodynamics. Your answer should include

- which of both quantities is cutoff-dependent,
- their relation to the wavefunction renormalization  $Z_2$  and the electron self-energy.

Also sketch how you would calculate  $Z_2$  in perturbation theory.

Only the physical mass  $m$  is an observable; it is the rest mass of an electron as measured in experiments. By contrast  $m_0$ , the so-called bare mass, is merely a parameter that specifies the Lagrangian. It can never be measured directly. Rather the Lagrangian produces measurable quantities, the scattering amplitudes, which depend on  $m_0$ .

This fact allows us to absorb the divergent quantity  $\delta m := m - m_0$  completely into  $m_0$ , making the latter a function of the measured physical mass and the cutoff  $\Lambda$ ,  $m_0 = m_0(m, \Lambda)$ .

The physical mass is the pole of the full propagator  $\frac{i}{\not{p} - m_0 - \Sigma(\not{p})}$ , i.e.  $m - m_0 = \Sigma(\not{p} = m)$ , where  $\Sigma$  is the 1PI electron self-energy (to order  $\alpha$  the one-loop photon exchange diagram), and the wavefunction renormalization  $Z_2$  is defined as

$$Z_2 := \left( 1 - \frac{d\Sigma(\not{p})}{d\not{p}} \right)^{-1} \Big|_{\not{p}=m}.$$

b) Draw all Feynman diagrams in QED which contribute to vertex renormalization at order  $e_0^4$  exactly.

This topic was unaddressed by our coverage of QFT. Part c) likewise.